



UWS Academic Portal

Enhanced airway smooth muscle relaxation via proteinase activated receptor 2

Black, K.; MacKenzie, A.; Dunning, L.; Crilly, A.; Brzeszczynska, J.; McGarvey, L.; Thornbury, K.; Goodyear, C.S.; Lockhart, J.C.; Litherland, G.J.

Published in:
Irish Journal of Medical Science

DOI:
[10.1007/s11845-019-02123-3](https://doi.org/10.1007/s11845-019-02123-3)

Published: 22/11/2019

Document Version
Peer reviewed version

[Link to publication on the UWS Academic Portal](#)

Citation for published version (APA):

Black, K., MacKenzie, A., Dunning, L., Crilly, A., Brzeszczynska, J., McGarvey, L., Thornbury, K., Goodyear, C. S., Lockhart, J. C., & Litherland, G. J. (2019). Enhanced airway smooth muscle relaxation via proteinase activated receptor 2. *Irish Journal of Medical Science*, 188(SUPPL 10), 286-286. <https://doi.org/10.1007/s11845-019-02123-3>

General rights

Copyright and moral rights for the publications made accessible in the UWS Academic Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact pure@uws.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

This is a post-peer-review, pre-copyedit version of an article published in Irish Journal of Medical Science. The final authenticated version is available online at: <http://dx.doi.org/10.1007/s11845-019-02123-3>

Enhanced airway smooth muscle relaxation via proteinase activated receptor 2

K. Black¹, A. MacKenzie¹, L. Dunning¹, A. Crilly¹, J. Brzezczynska¹, L. McGarvey², K. Thornbury³, C.S. Goodyear⁴, J.C. Lockhart¹, G.J. Litherland¹

¹Border & REgions Airways Training Hub, Institute of Biomedical and Environmental Health Research, School of Health & Life Sciences, University of the West of Scotland, Paisley, Scotland. ²Queens University Belfast, Belfast, Northern Ireland. ³Dundalk Institute of Technology, Dundalk, Ireland. ⁴Institute of Infection, Immunity & Inflammation, University of Glasgow.

Hyper-reactivity, inflammation and hyperplasia/hypertrophy of airway smooth muscle (ASM) limit airflow and are key features of chronic obstructive pulmonary disease (COPD). Proteinase activated receptor 2 (PAR2) is a critical modulator of inflammatory responses in respiratory disease such as asthma, yet is reported to promote ASM relaxation. However, the role of ASM PAR2 in COPD is not well understood¹.

Our aim was to determine the presence and role of PAR2 in murine lung and ASM subjected to oxidative stress using both immunohistochemistry (IHC) and wire myography.

PAR2 was detected using APR-32 antibody (Alomone, Israel) on both murine airway and lung tissue, with clean isotype.

Oxidative stress increased trypsin-induced ASM relaxation in both bronchial and tracheal tissue. This was PAR2 dependent, as relaxation was significantly reduced in PAR2^{-/-} compared with WT tracheal (mean \pm SEM; WT 53.4 \pm 13.4 % vs. PAR2^{-/-} 2.5 \pm 0.8 %; $p=0.02$; $n=3-4$) and bronchial tissue (WT 56 \pm 13.3 % vs. PAR2^{-/-} 2.3 \pm 0.9 %; $p=0.02$; $n=3-4$).

In conclusion, this study confirms a functional PAR2 role in murine airway tissue; importantly, the role of PAR2 in mediating ASM relaxation appears to be enhanced in oxidative environments such as found in COPD. This may have important implications for future potential therapies.

1. Sokolova E, Reiser G. A novel therapeutic target in various lung diseases: Airway proteases and protease-activated receptors. *Pharmacol Ther.* 2007;115(1):70–83.

This study was funded by the EU under the Interreg VA Programme, managed by the Special EU programmes body (SEUPB)